

## Lecture 26.

### Radiative and radiative-convective equilibrium.

#### Objectives:

1. Radiative equilibrium models.
2. Radiative-convective equilibrium models.

*Appendix.* Derivation of the Eddington gray radiative equilibrium.

#### Required reading:

L02: 8.3

### 1. Radiative equilibrium models.

These models predict the atmosphere temperature profile of an atmosphere in radiative

equilibrium  $\frac{dF_{net}}{dz} = 0$

Under the “gray atmosphere” assumption, we can solve for the temperature profile analytically

[\*Eddington gray radiative equilibrium results:\*](#) (see Appendix for the full derivation).

Assumptions:

- 1) Radiative equilibrium:  $\frac{dF_{net}}{dz} = 0$
- 2) Gray atmosphere in longwave
- 3) No scattering and black surface in longwave
- 4) No solar absorption in the atmosphere
- 5) Eddington approximation:  $I(\mu) = I_0 + I_1\mu$

Longwave flux profile:

$$F^\uparrow(\tau) = F_{sun} \left(1 + \frac{3}{4}\tau\right) \quad \text{and} \quad F^\downarrow(\tau) = F_{sun} \left(\frac{3}{4}\tau\right) \quad [26.1]$$

where  $F_{sun} = (1 - \bar{r})F_0 / 4$

Atmosphere blackbody emission and temperature profiles:

$$B(\tau) = \frac{F_{sun}}{2\pi} \left(1 + \frac{3}{2}\tau\right) \quad \text{and} \quad T^4(\tau) = T_e^4 \left(\frac{1}{2} + \frac{3}{4}\tau\right) \quad [26.2]$$

Surface temperature is discontinuous with the atmosphere (hotter):

$$B_s = B(\tau^*) + \frac{F_{sun}}{2\pi} \quad \text{and} \quad T_s^4 = T_e^4 \left(1 + \frac{3}{4}\tau^*\right) \quad [26.3]$$

Implications:

- ✓ Greenhouse effect – larger  $\tau^*$  increases surface temperature
- ✓ Runaway greenhouse effect -  $\tau^*$  increases  $\Rightarrow T_s$  increases
- ✓ Positive feedback – higher temperature  $\Rightarrow$  greenhouse gases

**Eddington gray radiative equilibrium temperatures**

If one wants the temperature profile in terms of height, one needs to relate optical depth to height.

Assume that an absorber has the exponential profile

$$\rho_a = \rho_0 \exp(-z/H_a) \quad [26.4]$$

So the profile of optical depth is

$$\tau(z) = \bar{k}_a \int_z^\infty \rho_a(z) dz = \bar{k}_a \rho_0 H_a \exp(-z/H_a) = \tau^* \exp(-z/H_a) \quad [26.5]$$

Temperature profile

$$T^4(z) = T_e^4 \left(1 + \frac{3}{4}\tau^* \exp(-z/H_a)\right) \quad [26.6]$$

Lapse rate

$$\frac{dT}{dz}(z) = -\frac{3}{8} \frac{\tau^*}{1 + \frac{3}{2}\tau^*} \frac{T(z)}{H_a} \exp(-z/H_a) \quad [26.7]$$

Implications:

- ✓ Low  $\tau^*$   $\Rightarrow$  stable atmosphere
- ✓ Smaller scale height  $H_a$  of the absorber causes steeper lapse rate
- ✓ Steepest lapse rate near the surface ( $z=0$ )

## 2. Radiative-convective equilibrium models.

- These climate models solve for the vertical profile of temperature using accurate broadband radiative transfer models.
- Model inputs vertical profile of gases, aerosols and clouds. Iterates the temperature profile to achieve equilibrium (i.e., zero heating rates or  $\frac{dF_{net}}{dz} = 0$ )
- Climate feedbacks can be included by having water vapor, surface albedo, clouds, etc. depend on temperature.

### Solving for radiative equilibrium:

Iterate the temperature profile  $T(z)$  to get zero heating rates  $\frac{\partial T}{\partial t} = 0$

#### 1. *Time marching method:*

$T$  at  $t+1$  time step from heating rate at time  $t$ :


$$T^{t+1}(z_k) = T(z_k) + \left( \frac{\partial T(z_k)}{\partial t} \right)^t \Delta t \quad [26.8]$$

#### 2. *Direct solve:*

Use gradient information in nonlinear root solver (faster, but more complex than time marching)

*Radiative equilibrium temperature profiles show (see Figure 26.1 below):*

- ✓ CO<sub>2</sub> –only-atmosphere has less steep profile.
- ✓ Earth's stratosphere warms due to UV absorption by ozone.
- ✓ Most greenhouse effect from water vapor.



**Figure 26.1** Pure radiative equilibrium temperature profiles for various atmospheric gases in a clear sky at 35 N in April. L+S means that the effects of both longwave and shortwave radiation are included (from Manabe and Strickler, 1964).

**Results:** the radiative equilibrium surface temperature is too high and the temperature profile is unrealistic.

**Problem:** radiative equilibrium surface temperature lapse rate near the surface exceeds threshold for convection

**Fix:** assume convection limits lapse rates to  $< \gamma_c$  (e.g., 6.5 K/km)


✓ **Radiative-convective equilibrium is equilibrium of radiative+ convective fluxes**

Convective adjustment methods:


1. Move heat like convection: if  $\gamma_c$  exceeded, adjust temperature so  $\gamma_c$  achieved and heat is conserved
2. Parameterize convective flux, e.g.

$$F_{conv} = C \left( \left| \frac{dT}{dz} \right| - \gamma_c \right) \quad \text{if} \left( \left| \frac{dT}{dz} \right| - \gamma_c \right) > 0 \quad [26.9]$$

*Results of the RCE model developed by Manabe and Strickler (1964):*



**Figure 26.2** Pure radiative equilibrium and radiative-convective equilibrium temperature profiles for two values of  $\gamma_c$  for clear sky.



**Figure 26.3** Radiative-convective equilibrium temperature profiles for various atmospheric gases in a clear sky at 35 N in April.

Comparing figures 26.1 and 26.3:

- ✓ Radiative equilibrium is fairly accurate for the stratosphere (though latitudinal and seasonal dependence is not correct)
- ✓ Convection required for reasonable tropospheric temperatures

## Appendix. Derivation of the Eddington gray radiative equilibrium.

Assumptions:

- 6) Radiative equilibrium:  $\frac{dF_{net}}{dz} = 0$
- 7) Gray atmosphere in longwave
- 8) No scattering and black surface in longwave
- 9) No solar absorption in the atmosphere
- 10) Eddington approximation:  $I(\mu) = I_0 + I_1\mu$

Since the atmosphere is gray (all wavelength are equivalent), one can write the wavelength integrated thermal emission radiative transfer equation ( no scattering)

$$\mu \frac{dI}{d\tau} = I - B$$

where  $I$  is the integrated radiance ( $\text{W m}^{-2} \text{sr}^{-1}$ ),  $\tau$  increases downward , and  $\mu > 0$  in the upward direction. Note that deriving the variation of  $B$  with the optical depth  $\tau$  is equivalent to determining the temperature profiles since the blackbody emission is a function of temperature only.

Using the Eddington approximation, the net flux (positive upward) becomes

$$F_{net} = 2\pi \int_{-1}^1 I\mu d\mu = \frac{4\pi}{3} I_1$$

The radiative equilibrium assumption implies that  $F_{net}$  (and  $I_1$ ) is constant with optical depth.

Integrating the above radiative transfer equation over  $d\mu$  gives

$$2\pi \frac{d}{d\tau} \int_{-1}^1 I\mu d\mu = 2\pi \int_{-1}^1 Id\mu - 2\pi \int_{-1}^1 Bd\mu$$
$$\frac{dF_{net}}{d\tau} = 4\pi I_0 - 4\pi B$$

Under the radiative equilibrium assumption, we have

$$I_0 = B$$

Integrating the radiative transfer equation over  $\mu d\mu$  gives

$$2\pi \frac{d}{d\tau} \int_{-1}^1 I \mu^2 d\mu = 2\pi \int_{-1}^1 I \mu d\mu - 2\pi \int_{-1}^1 B \mu d\mu$$

Since B is isotropic the last term drops out leaving

$$\frac{4\pi}{3} \frac{dI_0}{d\tau} = F_{net} = \frac{4\pi}{3} I_1$$

$$\frac{dB}{d\tau} = I_1$$

Thus, the solution for B is simply a linear function of optical depth:

$$B(\tau) = B(0) + I_1 \tau$$

Constants B(0) and  $I_1$  need to be determined from the boundary conditions.

Top of the atmosphere:

*First boundary condition:* no thermal downwelling flux

$$F^\downarrow(0) = 2\pi \int_{-1}^0 I \mu d\mu = \pi B(0) - \frac{2\pi}{3} I_1 = 0$$

so we have

$$I_1 = \frac{3}{2} B(0) \quad \text{or} \quad F_{net} = 2\pi B(0)$$

*Second boundary condition:* upwelling longwave flux is equal to the absorbed solar flux  $F_{sun}$ :

$$F^\uparrow(0) = 2\pi \int_0^1 I \mu d\mu = \pi B(0) + \frac{2\pi}{3} I_1 = F_{sun}$$

(Recall that the absorbed solar flux  $F_{sun}$  is  $F_{sun} = (1 - \bar{r}) F_0 / 4$  ) )

Putting in  $I_1 = \frac{3}{2} B(0)$  gives

$$F_{sun} = 2\pi B(0) = F_{net}$$

So now we have the B(0) and  $I_1$  and thus the atmosphere Planck function profile is determined

$$B(\tau) = \frac{F_{sun}}{2\pi} \left(1 + \frac{3}{2} \tau\right)$$

The final step is to apply the boundary condition at the surface to obtain the surface temperature  $T_s$ . This boundary condition is that the emitted flux by the surface equals to the sum of the downwelling shortwave and longwave flux at the black surface:



$$F_{sun} + F^{\downarrow}(\tau^*) = \pi B_s$$

where  $F^{\downarrow}(\tau^*) = \pi B(\tau^*) - \frac{2\pi}{3} I_1$

Using  $F_{sun} = \frac{4\pi}{3} I_1$  gives the emission from the surface

$$B_s = B(\tau^*) + \frac{F_{sun}}{2\pi}$$

which is discontinuous with the atmospheric emission.

The previous results can be expressed in terms of temperature by

$$T^4(\tau) = T_e^4 \left( \frac{1}{2} + \frac{3}{4} \tau \right)$$

where  $\sigma T_e^4 = F_{sun}$

$$T_{top}^4 = \frac{1}{2} T_e^4$$

$$T_s^4 = T_e^4 \left( 1 + \frac{3}{4} \tau^* \right)$$

For  $F_0 = 1366 \text{ W/m}^2$  and  $\bar{r} = 0.3$ :

$T_e = 255 \text{ K}$  and a “top” temperature  $T_t = 214 \text{ K}$

Assuming a global averaged surface air temperature of  $T(\tau^*) = 288 \text{ K}$  gives a gray body optical depth of  $\tau^* = 1.5$ , and a surface skin temperature of  $T_s = 308 \text{ K}$